


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 498/04, A61K 31/505 // (C07D 498/04, 261:00, 239:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/11003 (43) International Publication Date: 2 March 2000 (02.03.00)</p>
<p>(21) International Application Number: PCT/US99/18707 (22) International Filing Date: 18 August 1999 (18.08.99) (30) Priority Data: 60/097,386 21 August 1998 (21.08.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US], 974 Centre Road, WR-1ST18, Wilmington, DE 19807 (US). (72) Inventor: FRIETZE, William, Eric; 900 Merrybell Lane, Kennett Square, PA 19348 (US). (74) Agent: BROWDER, Monte, R.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).</p>		<p>(81) Designated States: AL, AU, BR, CA, CN, CZ, EE, HU, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.</p>
<p>(54) Title: ISOXAZOLO[4,5-d]PYRIMIDINES AS CRF ANTAGONISTS</p> <div style="text-align: center;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>The present invention describes novel isoxazolo[1,5-a]pyrimidines of formula (I), wherein R is an aromatic or heteroaromatic ring, or pharmaceutically acceptable salt form thereof, which are useful as CRF antagonists.</p>		

EZ

7

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AI.	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

ISOXAZOLO[4,5-d]PYRIMIDINES AS CRF ANTAGONISTS

5

FIELD OF THE INVENTION

This invention relates to novel isoxazolo[4,5-d]pyrimidines, pharmaceutical compositions containing the same and methods of using same in the treatment of psychiatric disorders and neurological diseases including affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing corticotropin releasing factor (CRF), including but not limited to disorders induced or facilitated by CRF.

25

BACKGROUND OF THE INVENTION

Corticotropin releasing factor, a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC)-derived peptide secretion from the anterior pituitary gland [J. Rivier et al., *Proc. Nat. Acad. Sci. (USA)* 80:4851 (1983); W. Vale et al., *Science* 213:1394 (1981)]. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone has a broad extrahypothalamic distribution in the central nervous system and produces a wide spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain [W. Vale et al., *Rec. Prog. Horm. Res.* 39:245 (1983); G.F. Koob, *Persp. Behav. Med.* 2:39 (1985); E.B. De Souza et al., *J. Neurosci.* 5:3189 (1985)].

There is also evidence that CRF plays a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors [J.E. Blalock, *Physiological Reviews* 69:1 (1989); J.E. Morley, *Life Sci.* 41:527 (1987)].

Clinical data provides evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system [for review see E.B. De Souza, *Hosp. Practice* 23:59 (1988)].

In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebral spinal fluid (CSF) of drug-free individuals [C.B. Nemeroff et al., *Science* 226:1342 (1984); C.M. Banki et al., *Am. J. Psychiatry* 144:873 (1987); R.D. France et al., *Biol. Psychiatry* 28:86 (1988); M. Arato et al., *Biol Psychiatry* 25:355 (1989)]. Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF [C.B. Nemeroff et al., *Arch. Gen. Psychiatry* 45:577 (1988)]. In addition, there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients [P.W. Gold et al., *Am J. Psychiatry* 141:619 (1984); F. Holsboer et al., *Psychoneuroendocrinology* 9:147 (1984); P.W. Gold et al., *New Eng. J. Med.* 314:1129 (1986)].

Preclinical studies in rats and non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression [R.M. Sapolsky, *Arch. Gen. Psychiatry* 46:1047 (1989)].


There is preliminary evidence that tricyclic antidepressants can alter CRF levels and thus modulate the numbers of CRF receptors in brain [Grigoriadis et al., *Neuropsychopharmacology* 2:53 (1989)].

There has also been a role postulated for CRF in the etiology of anxiety-related disorders. CRF produces anxiogenic effects in animals and interactions between benzodiazepine/non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models [D.R. Britton et al., *Life Sci.* 31:363 (1982); C.W. Berridge and A.J. Dunn *Regul. Peptides* 16:83 (1986)]. Preliminary studies using the putative CRF receptor antagonist α -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines [C.W. Berridge and A.J. Dunn, *Horm. Behav.* 21:393 (1987), *Brain Research Reviews* 15:71 (1990)]. Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics providing further evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic" effects of CRF in both the conflict test [K.T. Britton et al., *Psychopharmacology* 86:170 (1985); K.T. Britton et al., *Psychopharmacology* 94:306 (1988)] and in the acoustic startle test [N.R. Swerdlow et al., *Psychopharmacology* 88:147 (1986)] in rats. The benzodiazepine receptor antagonist (Ro15-1788), which was without behavioral activity alone in the operant conflict test, reversed the effects of CRF in a dose-dependent manner while the benzodiazepine inverse agonist (FG7142) enhanced the actions of CRF [K.T. Britton et al., *Psychopharmacology* 94:306 (1988)].

The mechanisms and sites of action through which the standard anxiolytics and antidepressants produce their therapeutic effects remain to be elucidated. It has been hypothesized however, that they are involved in the suppression of the CRF hypersecretion that is observed in these disorders. Of particular interest is that preliminary studies examining the effects of a CRF receptor antagonist (α -helical CRF₉₋₄₁) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the


benzodiazepines [for review see G.F. Koob and K.T. Britton, In: *Corticotropin-Releasing Factor: Basic and Clinical Studies of a Neuropeptide*, E.B. De Souza and C.B. Nemeroff eds., CRC Press p221 (1990)].

- 5 EP 0,778,277 (EP '277) describes a variety of CRF antagonists including those of the formula:




- 10 wherein A can be N or C, D can be N or C, E can be N or C, F can be N, C, S, or O, and G can be absent or a variety of groups. EP '277 does not disclose any isoxazolo[4,5-d]pyrimidines like those of the present invention.

- 15 WO 98/08847 (WO '847) describes a variety of CRF antagonists including those of the formula:



- 20 wherein A can be N or C, R⁵ is an aromatic or heteroaromatic group, and R and R³ can be a variety of groups. WO '847 does not disclose any isoxazolo[4,5-d]pyrimidines like those of the present invention.

- 25 WO 95/19774 (WO '774) describes a variety of CRF antagonists including those of the formula:



5 wherein A-B can be O, D can be N, and E can be C. However, WO '774 does not disclose any isoxazolo[4,5-d]pyrimidines like those of the present invention.

In view of the above, efficacious and specific antagonists of CRF are desired as potentially valuable
 10 therapeutic agents for the treatment of psychiatric disorders and neurological diseases. It is thus desirable to discover new CRF antagonists.


SUMMARY OF THE INVENTION

15 Accordingly, one object of the present invention is to provide novel isoxazolo[4,5-d]pyrimidines which are useful as CRF antagonists or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to
 20 provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt form thereof.

25 It is another object of the present invention to provide a method for treating psychiatric disorders and neurological diseases comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a
 30 pharmaceutically acceptable salt form thereof.

These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of formula I:




I

or pharmaceutically acceptable salt forms thereof, wherein R, R¹, and R² are defined below, are CRF antagonists.

DETAILED DESCRIPTION OF THE INVENTION

[1] A compound of formula I:



I

or a stereoisomer or pharmaceutically acceptable salt form thereof, wherein:

R is selected from phenyl, naphthyl, pyridyl, pyrimidinyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzthiazolyl, isoxazolyl and pyrazolyl, and is substituted with 0-5 R³ groups;

R¹ is selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, halo, -CN, C₁₋₄ haloalkyl, NR⁶R⁷, NR⁸COR⁹, -OR¹⁰, SH and -S(O)_nR¹¹;

R² is selected from NR^{6a}R^{7a} and OR^{10a};

R³ is independently selected at each occurrence from
 C₁₋₁₀ alkyl substituted with 0-2 R^a, C₂₋₁₀ alkenyl
 5 substituted with 0-2 R^a, C₂₋₁₀ alkynyl substituted with
 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a,
 (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2 R^a, -
 NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -
 NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -NR⁸CONR⁶R⁷, -
 10 C(O)OR^{10a}, SH, and -S(O)_nR¹²;

R^a is independently selected at each occurrence from
 C₁₋₄ alkyl, -NO₂, halo, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -
 NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -NR⁸CONR⁶R⁷, -
 15 C(O)OR^{10a}, SH, and -S(O)_nR¹²;

R⁶ is independently at each occurrence selected from H, C₁₋₄
 alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

20 R^{6a} is independently selected at each occurrence from H,
 C₁₋₆ alkyl substituted with 0-2 R^b, C₂₋₆ alkenyl
 substituted with 0-2 R^b, C₂₋₆ alkynyl substituted with
 0-2 R^b, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl substituted with
 0-2 R^b, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2
 25 R^b, aryl substituted with 0-2 R^b, (aryl)C₁₋₄ alkyl
 substituted with 0-2 R^b, heteroaryl substituted with
 0-2 R^b, (heteroaryl)C₁₋₄ alkyl substituted with 0-2 R^b,
 heterocyclyl substituted with 0-2 R^b, and
 (heterocyclyl)C₁₋₄ alkyl substituted with 0-2 R^b;

30 R^b is independently selected at each occurrence from C₁₋₄
 alkyl, -NO₂, halo, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -
 NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6b}R^{7a}, -NR⁸CONR^{6b}R^{7a}, -
 C(O)OR^{10a}, SH, and -S(O)_nR¹²;

35 R^{6b} is independently at each occurrence selected from H, C₁₋₄
 alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

R⁷ is independently at each occurrence selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

5 R^{7a} is independently selected at each occurrence from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)methyl, and C₁₋₄ haloalkyl;

alternatively, NR^{6a}R^{7a} is selected from piperidine, pyrrolidine, morpholine, thiomorpholine, 10 thiomorpholine-oxide, and thiomorpholine-dioxide, and is substituted with 0-1 R^e;

R^e is C₁₋₄ alkyl;

15 alternatively, NR^{6a}R^{7a} is piperazine or N-methylpiperazine, and is substituted with 0-1 R^f;

R^f is selected from C₁₋₄ alkyl, C(O)C₁₋₄ alkyl, C(O)benzyl, C(O)OC₁₋₄ alkyl, C(O)O-benzyl, SO₂-C₁₋₄ alkyl, SO₂- 20 benzyl, and SO₂-phenyl;

R⁸ is independently at each occurrence selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

25 R⁹ is independently at each occurrence selected from H, C₁₋₄ alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

R¹⁰ is independently at each occurrence selected from H, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, and 30 (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

R^{10a} is independently selected at each occurrence from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, and 35 C₁₋₄ haloalkyl;

R^{10b} is independently selected at each occurrence from H, C₁₋₆ alkyl, C₂₋₄ alkenyl, phenyl, benzyl, C₃₋₆

cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, and C₁₋₄ haloalkyl;

5 R¹¹ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, and C₃₋₆ cycloalkyl;

10 R¹² is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, aryl, and (aryl)C₁₋₄ alkyl;

15 R¹³ is independently at each occurrence selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

R^{13a} is independently at each occurrence selected from C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

20 aryl is independently at each occurrence selected from phenyl and naphthyl, and is substituted with 0-3 R⁹;

25 R⁹ is independently at each occurrence selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halo, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, SH, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -OC(O)R¹⁷, -NR⁸COR^{16a}, -NR⁸CONR^{6a}R^{7a}, -NR⁸C(O)OR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};

30 R¹⁵ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl, benzyl, (phenyl)C₁₋₄ alkyl and (naphthyl)C₁₋₄ alkyl;

35 R¹⁶ is independently at each occurrence selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

R^{16a} is independently at each occurrence selected from C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

5 R¹⁷ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl, benzyl, (phenyl)C₁₋₄ alkyl, (naphthyl)C₁₋₄ alkyl, heteroaryl and (heteroaryl)C₁₋₄ alkyl;

10

heteroaryl is independently at each occurrence selected from pyridyl, pyrimidinyl, triazinyl, furanyl, quinolinyl, isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, and indazolyl, and is substituted with 0-3 R^h;

15

heterocyclyl is saturated or partially saturated heteroaryl, substituted with 0-3 R^h; and

20

R^h is independently at each occurrence selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halo, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, SH, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -OC(O)R¹⁸, -NR⁸COR^{16a}, -NR⁸CONR^{6a}R^{7a}, -NR⁸CO₂R^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};

25

R¹⁸ is independently selected at each occurrence from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, phenyl and benzyl; and

30

n is independently at each occurrence selected from 0, 1 and 2.

35 [2] In a preferred embodiment, the present invention provides novel compounds, wherein:

R is selected from phenyl and pyridyl, and is substituted with 0-5 R³ groups;

5 R¹ is selected from H, CH₃, CH₂CH₃, Cl, F, Br, -CN, CF₃, NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, SH, SCH₃, and S(O)₂CH₃;

R² is NR^{6a}R^{7a}; and,

10 R³ is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, -
15 NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹².

20 [3] In a more preferred embodiment, the present invention provides novel compounds, wherein:

R is selected from phenyl and pyridyl, and is substituted with 1-3 R³ groups;

25 R¹ is selected from H, CH₃, CH₂CH₃, Cl, and F;

30 R³ is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, F, Cl, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR¹⁰, and -S(O)_nR¹².

35 R^a is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, F, Cl, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6a}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²;

- R^{6a} is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^b, C₂₋₆ alkenyl substituted with 0-2 R^b, C₂₋₆ alkynyl substituted with 0-2 R^b, C₃₋₆ cycloalkyl substituted with 0-2 R^b, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2 R^b, (aryl)C₁₋₄ alkyl substituted with 0-2 R^b, (heteroaryl)C₁₋₄ alkyl substituted with 0-2 R^b, and (heterocyclyl)C₁₋₄ alkyl substituted with 0-2 R^b;
- 10 R^b is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, F, Cl, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6b}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²; and,
- 15 R^{7a} is independently selected at each occurrence from H, C₁₋₆ alkyl, benzyl, and (C₃₋₆ cycloalkyl)methyl.
- [4] In an even more preferred embodiment, the present invention provides novel compounds, wherein:
- 20 R¹ is CH₃;
- R³ is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, -NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹²;
- 25 30 R^a is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, F, Cl, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹²;
- 35 R^{6a} is independently selected at each occurrence from C₂₋₄ alkyl substituted with 0-2 R^b, (C₃₋₅ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 groups selected from CH₃O and CH₃CH₂O, (aryl)C₁₋₂ alkyl substituted with 0-2 R^b,

(heteroaryl)C₁₋₂ alkyl substituted with 0-2 R^b, and
(heterocyclyl)C₁₋₂ alkyl substituted with 0-2 R^b;

5 R^b is independently selected at each occurrence from C₁₋₄
alkyl, -NO₂, F, Cl, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -COR¹³, -
OR^{10a}, -CONR^{6b}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²;

10 R⁸ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

10 R⁹ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

15 R¹⁰ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, and
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

20 R^{10a} is independently selected at each occurrence from H,
C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl,
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl, (C₁₋₄ alkoxy)C₁₋₂ alkyl, and
C₁₋₄ haloalkyl;

25 R^{10b} is independently selected at each occurrence from H,
C₁₋₆ alkyl, C₂₋₄ alkenyl, phenyl, benzyl, C₃₋₆
cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl, (C₁₋₄ alkoxy)C₁₋₂
alkyl, and C₁₋₄ haloalkyl;

30 R¹¹ is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, and C₃₋₆ cycloalkyl;

30 R¹² is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₂ alkyl,
C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl, aryl, and
(aryl)C₁₋₂ alkyl;

35 R^{13a} is independently at each occurrence selected from C₁₋₆
alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

aryl is independently at each occurrence phenyl substituted with 0-3 R^g;

- 5 R^g is independently at each occurrence selected from C₁₋₄ alkyl, C₃₋₆ cycloalkyl, F, Cl, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -NR⁸COR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};
- 10 R¹⁵ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl, benzyl, and (phenyl)C₁₋₂ alkyl;
- 15 R¹⁶ is independently at each occurrence selected from H, C₁₋₄ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;
- R^{16a} is independently at each occurrence selected from C₁₋₄ 20 alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

heteroaryl is independently at each occurrence selected from pyridyl, pyrimidinyl, furanyl, imidazolyl, thiazolyl, 25 pyrrolyl, oxazolyl, isoxazolyl, pyrazolyl, triazolyl, and tetrazolyl, and is substituted with 0-3 R^h;

heterocyclyl is saturated or partially saturated heteroaryl, substituted with 0-3 R^h; and,

- 30 R^h is independently at each occurrence selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, F, Cl, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -NR⁸COR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a}.

35

[5] In a further preferred embodiment, the present invention provides a compound selected from the group:

- 3-(2,4-dichlorophenyl)-5-methyl-7-(3-pentyl)amino-isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dichlorophenyl)-5-methyl-7-(methoxybut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(1,3-bis(methoxy)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dichlorophenyl)-5-methyl-7-(di(methoxyethyl)amino)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-butyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-propyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-diethylamino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dichlorophenyl)-5-methyl-7-dipropylamino-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-propyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dichlorophenyl)-5-methyl-7-(N-butyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dichlorophenyl)-5-methyl-7-di(cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dichlorophenyl)-5-methyl-7-(N-propyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-butyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dichlorophenyl)-5-methyl-7-dicyclopropylamino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4-dichlorophenyl)-5-methyl-7-(N-cyclopropyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 55

- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpent-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dichlorophenyl)-5-methyl-7-((1,3-dicyclopropyl)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbutyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dichlorophenyl)-5-methyl-7-((1,2-dicyclopropyl)ethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dichlorophenyl)-5-methyl-7-(1-hydroxy-but-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(morpholino)-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dichlorophenyl)-5-methyl-7-(3-pentoxy)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(methoxybut-2-oxy)-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpropoxy)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(dicyclopropylmethoxy)-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dichlorophenyl)-5-methyl-7-(2-methoxy-1-cyclopropylethoxy)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(3-pentyl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4,6-trimethylphenyl)-5-methyl-7-(methoxybut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4,6-trimethylphenyl)-5-methyl-7-(1,3-bis(methoxy)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(di(methoxyethyl)amino)-isoxazolo[4,5-d]pyrimidine;
- 55

- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-butyl)amino-isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-propyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-diethylamino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4,6-trimethylphenyl)-5-methyl-7-dipropylamino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-propyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-butyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4,6-trimethylphenyl)-5-methyl-7-di(cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-propyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-butyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4,6-trimethylphenyl)-5-methyl-7-dicyclopropylamino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-cyclopropyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylbut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylpent-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4,6-trimethylphenyl)-5-methyl-7-((1,3-dicyclopropyl)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylpropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylbutyl)amino-isoxazolo[4,5-d]pyrimidine;
- 55

- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4,6-trimethylphenyl)-5-methyl-7-((1,2-dicyclopropyl)ethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(1-hydroxy-but-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4,6-trimethylphenyl)-5-methyl-7-(morpholino)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(3-pentoxy)-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4,6-trimethylphenyl)-5-methyl-7-(methoxybut-2-oxy)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylpropoxy)-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4,6-trimethylphenyl)-5-methyl-7-(dicyclopropylmethoxy)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(2-methoxy-1-cyclopropylethoxy)-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dimethylphenyl)-5-methyl-7-(3-pentyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(methoxybut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dimethylphenyl)-5-methyl-7-(1,3-bis(methoxy)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dimethylphenyl)-5-methyl-7-(di(methoxyethyl)amino)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-butyl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-propyl)amino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4-dimethylphenyl)-5-methyl-7-diethylamino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-dipropylamino-isoxazolo[4,5-d]pyrimidine;
- 55 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;

- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-propyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dimethylphenyl)-5-methyl-7-(N-butyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-di(cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dimethylphenyl)-5-methyl-7-(N-propyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-butyl-N-cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dimethylphenyl)-5-methyl-7-dicyclopropylamino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-cyclopropyl-N-cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpent-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-((1,3-dicyclopropyl)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbutyl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dimethylphenyl)-5-methyl-7-(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-((1,2-dicyclopropyl)ethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dimethylphenyl)-5-methyl-7-(1-hydroxy-but-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4-dimethylphenyl)-5-methyl-7-(morpholino)-isoxazolo[4,5-d]pyrimidine;
- 55 3-(2,4-dimethylphenyl)-5-methyl-7-(3-pentoxy)-isoxazolo[4,5-d]pyrimidine;

- 3-(2,4-dimethylphenyl)-5-methyl-7-(methoxybut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpropoxy)-
isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dimethylphenyl)-5-methyl-7-(dicyclopropylmethoxy)-
isoxazolo[4,5-d]pyrimidine; and,
- 15 3-(2,4-dimethylphenyl)-5-methyl-7-(2-methoxy-1-
cyclopropylethoxy)isoxazolo[4,5-d]pyrimidine;

or a pharmaceutically acceptable salt form thereof.

- 20 In another embodiment, the present invention provides
pharmaceutical compositions comprising a pharmaceutically
acceptable carrier and a therapeutically effective amount of
at least one of the compounds of the present invention or a
pharmaceutically acceptable salt form thereof.

25

- In another embodiment, the present invention provides a
method of treating psychiatric disorders and neurological
diseases including affective disorder, anxiety, depression,
30 headache, irritable bowel syndrome, post-traumatic stress
disorder, supranuclear palsy, immune suppression,
Alzheimer's disease, gastrointestinal diseases, anorexia
nervosa or other feeding disorder, drug addiction, drug or
alcohol withdrawal symptoms, inflammatory diseases,
35 cardiovascular or heart-related diseases, fertility
problems, human immunodeficiency virus infections,
hemorrhagic stress, obesity, infertility, head and spinal
cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral
sclerosis, hypoglycemia or a disorder the treatment of which
40 can be effected or facilitated by antagonizing CRF,
including but not limited to disorders induced or
facilitated by CRF, in a mammal, comprising: administering
to the mammal a therapeutically effective amount of a

compound of formula (I) or a pharmaceutically acceptable salt form thereof.

5 In another embodiment, the present invention provides intermediate compounds useful in preparation of the CRF antagonist compounds and processes for making those intermediates, as described in the following description and claims.

10

The CRF antagonist compounds provided by this invention and labelled derivatives thereof are also useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the CRF receptor.

DEFINITIONS

The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a

substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties.

The present invention is intended to include all
5 isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

10 When any variable (e.g., R^6) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^6 , then said group may optionally
15 be substituted with up to two R^6 groups and R^6 at each occurrence is selected independently from the definition of R^6 . Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

20 When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given
25 formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "alkyl" is intended to include both
30 branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. "Haloalkyl" is intended to include
35 both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example $-C_vF_w$ where $v = 1$ to 3 and $w = 1$ to $(2v+1)$). Examples of

haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl.

"Alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxo, and s-pentoxo. "Cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, or cyclopentyl. Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl and propenyl. "Alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as ethynyl and propynyl.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo; and "counter-ion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7-to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl.

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon

atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heterotams independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl,

phenanthridinyl, phenanthrolinyl, phenazinyl,
phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl,
piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl,
pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,
5 pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole,
pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl,
2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl,
4H-quinolizinyl, quinoxalinyl, quinuclidinyl,
tetrahydrofuranyl, tetrahydroisoquinolinyl,
10 tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-
thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-
thiadiazolyl, thianthrenyl, thiazolyl, thienyl,
thienothiazolyl, thienooxazolyl, thienoimidazolyl,
thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
15 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl. Preferred
heterocycles include, but are not limited to, pyridinyl,
furanyl, thienyl, pyrrolyl, pyrazolyl, pyrrolidinyl,
imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl,
oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl,
20 benzoxazolyl, and isatinoyl. Also included are fused ring
and spiro compounds containing, for example, the above
heterocycles.

The phrase "pharmaceutically acceptable" is employed
herein to refer to those compounds, materials, compositions,
25 and/or dosage forms which are, within the scope of sound
medical judgment, suitable for use in contact with the
tissues of human beings and animals without excessive
toxicity, irritation, allergic response, or other problem or
complication, commensurate with a reasonable benefit/risk
30 ratio.

As used herein, "pharmaceutically acceptable salts"
refer to derivatives of the disclosed compounds wherein the
parent compound is modified by making acid or base salts
thereof. Examples of pharmaceutically acceptable salts
35 include, but are not limited to, mineral or organic acid
salts of basic residues such as amines; alkali or organic
salts of acidic residues such as carboxylic acids; and the
like. The pharmaceutically acceptable salts include the

conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from

5 inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic,

10 phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which

15 contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two;

20 generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby

25 incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound

30 of formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula (I) wherein a hydroxy, amino, or sulfhydryl group is bonded

35 to any group that, when the prodrug or compound of formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not

limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula (I), and the like.

Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. A stable compound or stable structure is meant to imply a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an effective therapeutic agent.

The term "therapeutically effective amount" of a compound of this invention means an amount effective to antagonize abnormal level of CRF or treat the symptoms of affective disorder, anxiety or depression in a host.

Synthesis


The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. All references cited herein are hereby incorporated in their entirety herein by reference.

The following abbreviations are used herein:

AcOH	acetic acid
t-BuOK	potassium tert-butoxide
DMSO	dimethyl sulfoxide
EtOAc	ethyl acetate
EtOH	ethanol
THF	tetrahydrofuran
TLC	thin layer chromatography

The compounds of this invention may be prepared using the methods shown in Scheme 1. The synthesis proceeds via an α -nitroacetophenone (X) which may be prepared by C-acylation of nitromethane with aroyl phenyl esters

(Synthesis, 1979, p295) or with substituted phenyl carboxylic acids (Chem. Pharm. Bull., 1981, 29(1), p259). The ketone (X) is then converted to the oxime (XI) which is treated with an alkyl oxalyl chloride to give an oxime
5 oxalic ester. The ester is subjected to base promoted cyclization to yield isoxazole (XII) (Heterocycles, 1985, 23(6), p1465). The nitro group is then reduced using an appropriate method, for example, zinc in aqueous ammonium chloride, to produce aminoisoxazole (XIII). Amidation of
10 the ester (XIII) may be accomplished by treatment with ammonium hydroxide to yield amide (XIV), which can be cyclized using, for example, orthoesters, to yield isoxazolopyrimidones of the formula (XV). Conversion of the hydroxy group in the tautomeric form of (XV) to a suitable
15 leaving group may be accomplished, for example, by chlorination using, phosphorous oxychloride, or by sulfonylation (e.g., SO_2Cl_2 /collidine or $\text{CH}_3\text{SO}_2\text{Cl}/\text{Et}_3\text{N}$) to give a sulfonate as a leaving group. Treatment of (XVI) with amines in the presence of organic or inorganic bases
20 provides compounds of formula (I) where R^2 = substituted amino.



Analogous treatment of (XVI) with alcohols, optionally in the presence of bases such as NaH , provides compounds of Formula (I) where $\text{R}^2 = \text{e.g., alkoxy}$.

If intermediates contain functional groups which are sensitive to reaction conditions, these groups may be protected using methods known to those skilled in the art. These methods include, but are not limited to, those described in Protective Groups in Organic Synthesis (Greene, Wuts; 2nd ed., 1991, John Wiley & Sons, Inc.).

The following examples are provided to describe the invention in further detail. These examples, which set forth the best mode presently contemplated for carrying out the invention, are intended to illustrate and not to limit the invention.

Examples

Abbreviations used in the Examples are defined as follows: "1 x" for once, "2 x" for twice, "3 x" for thrice, "°C" for degrees Celsius, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "mL" for milliliter or milliliters, "¹H" for proton, "h" for hour or hours, "M" for molar, "min" for minute or minutes, "MHz" for megahertz, "MS" for mass spectroscopy, "NMR" for nuclear magnetic resonance spectroscopy, "rt" for room temperature, "tlc" for thin layer chromatography, "v/v" for volume to volume ratio. "α", "β", "R" and "S" are stereochemical designations familiar to those skilled in the art.

Example 1

3-(2,4-Dichlorophenyl)-5-methyl-7-(1-ethylpropyl)amino-isoxazolo[4,5-d]pyrimidine

Part A: To a cooled (ice bath) solution of 2,4-dichlorobenzoyl chloride (5mL, 35.66mmol) in 50mL CH₂Cl₂ was added phenol (3.13mL, 35.66mmol), followed by triethylamine (4.97mL, 35.66mmol). The resulting reaction mixture was stirred at room temperature for 5h. The mixture was then partitioned between CH₂Cl₂ and water. The organic layer was dried (Na₂SO₄), filtered, and evaporated in vacuo to yield 9.5g of phenyl 2,4-dichlorobenzoate. MS (NH₃-CI) 284 (M+NH₄)⁺.

Part B: To a suspension of t-BuOK (12g, 107mmol) in 160mL DMSO was cautiously added nitromethane (5.8mL, 107mmol) with stirring. After 1h, phenyl 2,4-dichlorobenzoate (9.5g, 35.6mmol), was slowly added and the resulting reaction

mixture was stirred at room temperature for 3h. 100mL of cold water and urea (3.3g) were added and the mixture was cooled in an ice bath then acidified using conc. HCl (10mL). The product was precipitated by pouring the mixture into 700mL of water. 7.1g of alpha-nitro-2,4-dichloroacetophenone was collected as a pale yellow solid. ^1H NMR (300MHz, CDCl_3) δ 7.78 (d, 1H), 7.53 (m, 1H), 7.43 (m, 1H), 5.9 (s, 2H).

10 Part C: A mixture of alpha-nitro-2,4-dichloroacetophenone (8.3g, 35.6mmol) and hydroxylamine hydrochloride (2.5g, 35.6mmol) in ethanol (30mL) and AcOH (10mL) was refluxed for 7h. The mixture was concentrated *in vacuo* and the residue was partitioned between EtOAc and water. The EtOAc layer was dried (Na_2SO_4), filtered, and evaporated *in vacuo* to yield 8.54g of crude oxime. The crude product was dissolved in Et_2O and ethyl oxalyl chloride (3.83mL, 34.29mmol) was added. The solution was stirred at room temperature for 16h, then small amounts of triethylamine were added as product formation was monitored by TLC. When the reaction was complete, the reaction mixture was partitioned between Et_2O and water. The organic layer was dried (Na_2SO_4), filtered, and evaporated *in vacuo* to yield a residue which was purified by column chromatography on silica gel (100% hexane to 50% EtOAc in hexane gradient) to yield 5.02g of the desired nitro isoxazole.

Part D: To a cooled (ice bath) mixture of the nitroisoxazole from Example 1, Part C (4.75g, 14.35mmol) and NH_4Cl (17.96g, 335.69mmol) in 75mL of water, was added zinc dust (7.98g, 121.98mmol) portionwise over 15 min. The mixture was stirred for 2h, then 20mL of EtOAc was added and the mixture was stirred for 45 min. The mixture was filtered through celite and partitioned between EtOAc and water. The organic layer was evaporated *in vacuo* to afford a residue corresponding to the aminoisoxazole. MS ($\text{NH}_3\text{-CI}$) 301 ($\text{M}+\text{H}$) $^+$; 318 ($\text{M}+\text{NH}_3$) $^+$. The residue was dissolved in THF (25mL). To this solution was added 100mL of ammonium

hydroxide and 10mL of methanol. The mixture was stoppered and stirred at room temperature for 16h. Volatiles were removed *in vacuo* and the concentrate was partitioned between EtOAc and water. The organic layer was dried (Na₂SO₄),
5 filtered, and evaporated *in vacuo* to yield the amide. MS (NH₃-CI) 272 (M+H)⁺; 289 (M+NH₃)⁺. This crude product was refluxed in triethyl orthoacetate (50 mL) for 3h, then concentrated *in vacuo*. The resulting residue was dissolved in 10mL THF. 20mL of 3M NaOH was added and the mixture was
10 refluxed for 45min, then concentrated *in vacuo*. The concentrate was diluted with water and AcOH was added to a pH between 5-6. This mixture was partitioned between EtOAc and water. The organic layer was evaporated *in vacuo* and the residue was purified by column chromatography on silica
15 gel to give 798mg of the desired isoxazolopyrimidone. MS (NH₃-CI) 296 (M+H)⁺; 313 (M+NH₃)⁺.

Part E: The isoxazolopyrimidone from Example 1, Part D (790mg) was refluxed in 10mL of POCl₃ for 3h. The reaction
20 mixture was concentrated *in vacuo* and the residue was purified by column chromatography on silica gel (100% hexane to 25% EtOAc in Hexane gradient) to give 568mg of the chloro isoxazolopyrimidine as a pale yellow solid. MS (NH₃-CI) 314 (M+H)⁺.

25 Part F: To a solution of 30mg (0.095 mmol) of the chloro isoxazolopyrimidine (Example 1, Part E) in methylene chloride (3mL) was added a solution of 9.15mg (0.105 mmol) of 1-ethylpropylamine in methylene chloride
30 (0.5mL), followed by 0.015mL (0.105 mmol) of triethylamine. The resulting solution was stirred at room temperature for 16 h. The reaction mixture was added to a column containing silica gel and eluted (100% hexane to 25% EtOAc in Hexane gradient) to give 26mg of the title compound as a colorless
35 viscous oil. MS (NH₃-CI) 365 (M+H)⁺. HRMS: Mass = 365.0938; Calc. Mass = 365.0936 for C₁₇H₁₉N₄OCl₂ (M+H)⁺. ¹H NMR (300MHz, CDCl₃) 8.7.82 (m, 1H), 7.6 (d, 1H, J = 2.2Hz), 7.43 (dd, 1H, J = 2.2, 6.22, 1.83Hz), 5.15 (bm, 1H), 4.32

WO 00/11003


PCT/US99/18707

(bm, 1H), 2.6 (s, 3H), 1.75 (m, 2H), 1.61 (m, 2H), 1.00 (t, 6H, J = 7.3Hz).

Compounds in Table 1 not previously described can be made by similar procedures.

Table 1

5



Ex.	R ¹	R ²	R	MS ^a
1	CH ₃	NHCH(CH ₂ CH ₃) ₂	2,4-Cl ₂ Ph	365
2	CH ₃	NHCH(CH ₂ OCH ₃)Et	2,4-Cl ₂ Ph	381
3	CH ₃	NHCH(CH ₂ OCH ₃) ₂	2,4-Cl ₂ Ph	397.1
4	CH ₃	N(CH ₂ CH ₂ OCH ₃) ₂	2,4-Cl ₂ Ph	411.1
5	CH ₃	N(Bu)Et	2,4-Cl ₂ Ph	379.1
6	CH ₃	N(Pr)Et	2,4-Cl ₂ Ph	
7	CH ₃	N(Et) ₂	2,4-Cl ₂ Ph	
8	CH ₃	N(Pr) ₂	2,4-Cl ₂ Ph	
9	CH ₃	N(CH ₂ C-Pr)Et	2,4-Cl ₂ Ph	
10	CH ₃	N(CH ₂ C-Pr)Pr	2,4-Cl ₂ Ph	
11	CH ₃	N(CH ₂ C-Pr)Bu	2,4-Cl ₂ Ph	
12	CH ₃	N(CH ₂ C-Pr) ₂	2,4-Cl ₂ Ph	
13	CH ₃	N(c-Pr)Et	2,4-Cl ₂ Ph	
14	CH ₃	N(c-Pr)Pr	2,4-Cl ₂ Ph	
15	CH ₃	N(c-Pr)Bu	2,4-Cl ₂ Ph	
16	CH ₃	N(c-Pr) ₂	2,4-Cl ₂ Ph	
17	CH ₃	N(CH ₂ C-Pr)c-Pr	2,4-Cl ₂ Ph	
18	CH ₃	NHCH(CH ₂ C-Pr)Et	2,4-Cl ₂ Ph	
19	CH ₃	NHCH(CH ₂ C-Pr)Pr	2,4-Cl ₂ Ph	
20	CH ₃	NHCH(CH ₂ C-Pr) ₂	2,4-Cl ₂ Ph	
21	CH ₃	NHCH(c-Pr)Et	2,4-Cl ₂ Ph	
22	CH ₃	NHCH(c-Pr)Pr	2,4-Cl ₂ Ph	
23	CH ₃	NHCH(c-Pr) ₂	2,4-Cl ₂ Ph	
24	CH ₃	NHCH(CH ₂ C-Pr)c-Pr	2,4-Cl ₂ Ph	
25	CH ₃	NHCH(CH ₂ OH)Et	2,4-Cl ₂ Ph	367
26	CH ₃	morpholino	2,4-Cl ₂ Ph	365

27	CH ₃	OCH(Et) ₂	2,4-Cl ₂ Ph	
28	CH ₃	OCH(CH ₂ OCH ₃)Et	2,4-Cl ₂ Ph	382
29	CH ₃	OCH(CH ₂ c-Pr)Et	2,4-Cl ₂ Ph	
30	CH ₃	OCH(c-Pr)Et	2,4-Cl ₂ Ph	
31	CH ₃	OCH(c-Pr) ₂	2,4-Cl ₂ Ph	
32	CH ₃	OCH(CH ₂ OCH ₃)c-Pr	2,4-Cl ₂ Ph	
33	CH ₃	NHCH(CH ₂ CH ₃) ₂	2,4-(CH ₃) ₂ Ph	325.2
34	CH ₃	NHCH(CH ₂ OCH ₃)Et	2,4-(CH ₃) ₂ Ph	341.2
35	CH ₃	NHCH(CH ₂ OCH ₃) ₂	2,4-(CH ₃) ₂ Ph	
36	CH ₃	N(CH ₂ CH ₂ OCH ₃) ₂	2,4-(CH ₃) ₂ Ph	371.2
37	CH ₃	N(Bu)Et	2,4-(CH ₃) ₂ Ph	
38	CH ₃	N(Pr)Et	2,4-(CH ₃) ₂ Ph	
39	CH ₃	N(Et) ₂	2,4-(CH ₃) ₂ Ph	
40	CH ₃	N(Pr) ₂	2,4-(CH ₃) ₂ Ph	
41	CH ₃	N(CH ₂ c-Pr)Et	2,4-(CH ₃) ₂ Ph	
42	CH ₃	N(CH ₂ c-Pr)Pr	2,4-(CH ₃) ₂ Ph	
43	CH ₃	N(CH ₂ c-Pr)Bu	2,4-(CH ₃) ₂ Ph	
44	CH ₃	N(CH ₂ c-Pr) ₂	2,4-(CH ₃) ₂ Ph	
45	CH ₃	N(c-Pr)Et	2,4-(CH ₃) ₂ Ph	
46	CH ₃	N(c-Pr)Pr	2,4-(CH ₃) ₂ Ph	
47	CH ₃	N(c-Pr)Bu	2,4-(CH ₃) ₂ Ph	
48	CH ₃	N(c-Pr) ₂	2,4-(CH ₃) ₂ Ph	
49	CH ₃	N(CH ₂ c-Pr)c-Pr	2,4-(CH ₃) ₂ Ph	
50	CH ₃	NHCH(CH ₂ c-Pr)Et	2,4-(CH ₃) ₂ Ph	
51	CH ₃	NHCH(CH ₂ c-Pr)Pr	2,4-(CH ₃) ₂ Ph	
52	CH ₃	NHCH(CH ₂ c-Pr) ₂	2,4-(CH ₃) ₂ Ph	
53	CH ₃	NHCH(c-Pr)Et	2,4-(CH ₃) ₂ Ph	
54	CH ₃	NHCH(c-Pr)Pr	2,4-(CH ₃) ₂ Ph	
55	CH ₃	NHCH(c-Pr) ₂	2,4-(CH ₃) ₂ Ph	
56	CH ₃	NHCH(CH ₂ c-Pr)c-Pr	2,4-(CH ₃) ₂ Ph	
57	CH ₃	NHCH(CH ₂ OH)Et	2,4-(CH ₃) ₂ Ph	
58	CH ₃	morpholino	2,4-(CH ₃) ₂ Ph	
59	CH ₃	OCH(Et) ₂	2,4-(CH ₃) ₂ Ph	
60	CH ₃	OCH(CH ₂ OCH ₃)Et	2,4-(CH ₃) ₂ Ph	
61	CH ₃	OCH(CH ₂ c-Pr)Et	2,4-(CH ₃) ₂ Ph	
62	CH ₃	OCH(c-Pr)Et	2,4-(CH ₃) ₂ Ph	
63	CH ₃	OCH(c-Pr) ₂	2,4-(CH ₃) ₂ Ph	

64	CH ₃	OCH(CH ₂ OCH ₃) c-Pr	2,4-(CH ₃) ₂ Ph
65	CH ₃	NHCH(CH ₂ CH ₃) ₂	2,4,6-(CH ₃) ₃ Ph
66	CH ₃	NHCH(CH ₂ OCH ₃) Et	2,4,6-(CH ₃) ₃ Ph
67	CH ₃	NHCH(CH ₂ OCH ₃) ₂	2,4,6-(CH ₃) ₃ Ph
68	CH ₃	N(CH ₂ CH ₂ OCH ₃) ₂	2,4,6-(CH ₃) ₃ Ph
69	CH ₃	N(Bu) Et	2,4,6-(CH ₃) ₃ Ph
70	CH ₃	N(Pr) Et	2,4,6-(CH ₃) ₃ Ph
71	CH ₃	N(Et) ₂	2,4,6-(CH ₃) ₃ Ph
72	CH ₃	N(Pr) ₂	2,4,6-(CH ₃) ₃ Ph
73	CH ₃	N(CH ₂ c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
74	CH ₃	N(CH ₂ c-Pr) Pr	2,4,6-(CH ₃) ₃ Ph
75	CH ₃	N(CH ₂ c-Pr) Bu	2,4,6-(CH ₃) ₃ Ph
76	CH ₃	N(CH ₂ c-Pr) ₂	2,4,6-(CH ₃) ₃ Ph
77	CH ₃	N(c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
78	CH ₃	N(c-Pr) Pr	2,4,6-(CH ₃) ₃ Ph
79	CH ₃	N(c-Pr) Bu	2,4,6-(CH ₃) ₃ Ph
80	CH ₃	N(c-Pr) ₂	2,4,6-(CH ₃) ₃ Ph
81	CH ₃	N(CH ₂ c-Pr) c-Pr	2,4,6-(CH ₃) ₃ Ph
82	CH ₃	NHCH(CH ₂ c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
83	CH ₃	NHCH(CH ₂ c-Pr) Pr	2,4,6-(CH ₃) ₃ Ph
84	CH ₃	NHCH(CH ₂ c-Pr) ₂	2,4,6-(CH ₃) ₃ Ph
85	CH ₃	NHCH(c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
86	CH ₃	NHCH(c-Pr) Pr	2,4,6-(CH ₃) ₃ Ph
87	CH ₃	NHCH(c-Pr) ₂	2,4,6-(CH ₃) ₃ Ph
88	CH ₃	NHCH(CH ₂ c-Pr) c-Pr	2,4,6-(CH ₃) ₃ Ph
89	CH ₃	NHCH(CH ₂ OH) Et	2,4,6-(CH ₃) ₃ Ph
90	CH ₃	morpholino	2,4,6-(CH ₃) ₃ Ph
91	CH ₃	OCH(Et) ₂	2,4,6-(CH ₃) ₃ Ph
92	CH ₃	OCH(CH ₂ OCH ₃) Et	2,4,6-(CH ₃) ₃ Ph
93	CH ₃	OCH(CH ₂ c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
94	CH ₃	OCH(c-Pr) Et	2,4,6-(CH ₃) ₃ Ph
95	CH ₃	OCH(c-Pr) ₂	2,4,6-(CH ₃) ₃ Ph
96	CH ₃	OCH(CH ₂ OCH ₃) c-Pr	2,4,6-(CH ₃) ₃ Ph

^a Observed (M+H)⁺ ion under (NH₃-CI) MS conditions.

Utility

Compounds of this invention are expected to have utility in the treatment of imbalances associated with abnormal levels of CRF in patients suffering from depression, affective disorders, and/or anxiety.

CRF-R1 Receptor Binding Assay for the Evaluation of Biological Activity

The following is a description of the isolation of cell membranes containing cloned human CRF-R1 receptors for use in the standard binding assay as well as a description of the assay itself.

Messenger RNA was isolated from human hippocampus. The mRNA was reverse transcribed using oligo (dt) 12-18 and the coding region was amplified by PCR from start to stop codons. The resulting PCR fragment was cloned into the EcoRV site of pGEMV, from whence the insert was reclaimed using XhoI + XbaI and cloned into the XhoI + XbaI sites of vector pm3ar (which contains a CMV promoter, the SV40 't' splice and early poly A signals, an Epstein-Barr viral origin of replication, and a hygromycin selectable marker). The resulting expression vector, called phchCRFR was transfected in 293EBNA cells and cells retaining the episome were selected in the presence of 400 mM hygromycin. Cells surviving 4 weeks of selection in hygromycin were pooled, adapted to growth in suspension and used to generate membranes for the binding assay described below. Individual aliquots containing approximately 1×10^8 of the suspended cells were then centrifuged to form a pellet and frozen.

For the binding assay a frozen pellet described above containing 293EBNA cells transfected with hCRFR1 receptors is homogenized in 10 ml of ice cold tissue buffer (50 mM HEPES buffer pH 7.0, containing 10 mM $MgCl_2$, 2 mM EGTA, 1 mg/ml aprotinin, 1 mg/ml leupeptin and 1 mg/ml pepstatin). The homogenate is centrifuged at 40,000 x g for 12 min and the resulting pellet rehomogenized in 10 ml of tissue buffer. After another centrifugation at 40,000 x g for 12

min, the pellet is resuspended to a protein concentration of 360 mg/ml to be used in the assay.

Binding assays are performed in 96 well plates; each well having a 300 ml capacity. To each well is added 50 ml of test drug dilutions (final concentration of drugs range from 10^{-10} - 10^{-5} M), 100 ml of ^{125}I -ovine-CRF (^{125}I -o-CRF) (final concentration 150 pM) and 150 ml of the cell homogenate described above. Plates are then allowed to incubate at room temperature for 2 hours before filtering the incubate over GF/F filters (presoaked with 0.3% polyethylenimine) using an appropriate cell harvester. Filters are rinsed 2 times with ice cold assay buffer before removing individual filters and assessing them for radioactivity on a gamma counter.

Curves of the inhibition of ^{125}I -o-CRF binding to cell membranes at various dilutions of test drug are analyzed by the iterative curve fitting program LIGAND [P.J. Munson and D. Rodbard, *Anal. Biochem.* 107:220 (1980)], which provides K_i values for inhibition which are then used to assess biological activity.

A compound is considered to be active if it has a K_i value of less than about 10000 nM for the inhibition of CRF.

Inhibition of CRF-Stimulated Adenylate Cyclase Activity

Inhibition of CRF-stimulated adenylate cyclase activity can be performed as described by G. Battaglia et al. *Synapse* 1:572 (1987). Briefly, assays are carried out at 37° C for 10 min in 200 ml of buffer containing 100 mM Tris-HCl (pH 7.4 at 37° C), 10 mM MgCl_2 , 0.4 mM EGTA, 0.1% BSA, 1 mM isobutylmethylxanthine (IBMX), 250 units/ml phosphocreatine kinase, 5 mM creatine phosphate, 100 mM guanosine 5'-triphosphate, 100 nM oCRF, antagonist peptides (concentration range 10^{-9} to 10^{-6}M) and 0.8 mg original wet weight tissue (approximately 40-60 mg protein). Reactions are initiated by the addition of 1 mM ATP/ ^{32}P ATP (approximately 2-4 mCi/tube) and terminated by the addition of 100 ml of 50 mM Tris-HCL, 45 mM ATP and 2% sodium dodecyl sulfate. In order to monitor the recovery of cAMP, 1 μl of

[³H]cAMP (approximately 40,000 dpm) is added to each tube prior to separation. The separation of [³²P]cAMP from [³²P]ATP is performed by sequential elution over Dowex and alumina columns.

5

In vivo Biological Assay

The *in vivo* activity of the compounds of the present invention can be assessed using any one of the biological assays available and accepted within the art. Illustrative of these tests include the Acoustic Startle Assay, the Stair Climbing Test, and the Chronic Administration Assay. These and other models useful for the testing of compounds of the present invention have been outlined in C.W. Berridge and A.J. Dunn *Brain Research Reviews* 15:71 (1990). Compounds may be tested in any species of rodent or small mammal.

15

Dosage and Formulation

Compounds of this invention can be administered to treat these abnormalities by means that produce contact of the active agent with the agent's site of action in the body of a mammal. The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals either as individual therapeutic agent or in combination of therapeutic agents. They can be administered alone, but will generally be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

25

The dosage administered will vary depending on the use and known factors such as pharmacodynamic character of the particular agent, and its mode and route of administration; the recipient's age, weight, and health; nature and extent of symptoms; kind of concurrent treatment; frequency of treatment; and desired effect. For use in the treatment of said diseases or conditions, the compounds of this invention can be orally administered daily at a dosage of the active ingredient of 0.002 to 200 mg/kg of body weight. Ordinarily, a dose of 0.01 to 10 mg/kg in divided doses one

30

35

to four times a day, or in sustained release formulation will be effective in obtaining the desired pharmacological effect.

Dosage forms (compositions) suitable for administration contain from about 1 mg to about 100 mg of active ingredient per unit. In these pharmaceutical compositions, the active ingredient will ordinarily be present in an amount of about 0.5 to 95% by weight based on the total weight of the composition.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets and powders; or in liquid forms such as elixirs, syrups, and/or suspensions. The compounds of this invention can also be administered parenterally in sterile liquid dose formulations.

Gelatin capsules can be used to contain the active ingredient and a suitable carrier such as but not limited to lactose, starch, magnesium stearate, steric acid, or cellulose derivatives. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of time. Compressed tablets can be sugar-coated or film-coated to mask any unpleasant taste, or used to protect the active ingredients from the atmosphere, or to allow selective disintegration of the tablet in the gastrointestinal tract.

Liquid dose forms for oral administration can contain coloring or flavoring agents to increase patient acceptance.

In general, water, pharmaceutically acceptable oils, saline, aqueous dextrose (glucose), and related sugar solutions and glycols, such as propylene glycol or polyethylene glycol, are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents, such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or in combination, are suitable stabilizing agents. Also used

are citric acid and its salts, and EDTA. In addition, parenteral solutions can contain preservatives such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

5 Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences", A. Osol, a standard reference in the field.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as
10 follows:

Capsules

A large number of units capsules are prepared by filling standard two-piece hard gelatin capsules each with 100 mg of powdered active ingredient, 150 mg lactose, 50 mg
15 cellulose, and 6 mg magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestible oil such as soybean, cottonseed oil, or olive oil is prepared and injected by means of a positive displacement was pumped into
20 gelatin to form soft gelatin capsules containing 100 mg of the active ingredient. The capsules were washed and dried.

Tablets


A large number of tablets are prepared by conventional procedures so that the dosage unit was 100 mg active
25 ingredient, 0.2 mg of colloidal silicon dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg of starch, and 98.8 mg lactose. Appropriate coatings may be applied to increase palatability or delayed adsorption.

The compounds of this invention may also be used as
30 reagents or standards in the biochemical study of neurological function, dysfunction, and disease.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the
35 scope of the appended claims, the invention may be practiced otherwise that as specifically described herein.

WHAT IS CLAIMED AS NEW AND DESIRED TO BE SECURED BY LETTER
PATENT OF UNITED STATES IS:

1. A compound of formula (I):



I

or a stereoisomer or pharmaceutically acceptable salt form thereof, wherein:

R is selected from phenyl, naphthyl, pyridyl, pyrimidinyl, triazinyl, furanyl, quinoliny, isoquinoliny, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuranyl, benzothienyl, benzthiazolyl, isoxazolyl and pyrazolyl, and is substituted with 0-5 R³ groups;

R¹ is selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, halo, -CN, C₁₋₄ haloalkyl, NR⁶R⁷, NR⁸COR⁹, -OR¹⁰, SH and -S(O)_nR¹¹;

R² is selected from NR^{6a}R^{7a} and OR^{10a};

R³ is independently selected at each occurrence from C₁₋₁₀ alkyl substituted with 0-2 R^a, C₂₋₁₀ alkenyl substituted with 0-2 R^a, C₂₋₁₀ alkynyl substituted with 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2 R^a, -NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -NR⁸CONR⁶R⁷, -C(O)OR^{10a}, SH, and -S(O)_nR¹²;

R^a is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, halo, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -

NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -NR⁸CONR⁶R⁷, -
C(O)OR^{10a}, SH, and -S(O)_nR¹²;

5 R⁶ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

10 R^{6a} is independently selected at each occurrence from H,
C₁₋₆ alkyl substituted with 0-2 R^b, C₂₋₆ alkenyl
substituted with 0-2 R^b, C₂₋₆ alkynyl substituted with
0-2 R^b, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl substituted with
0-2 R^b, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2
R^b, aryl substituted with 0-2 R^b, (aryl)C₁₋₄ alkyl
substituted with 0-2 R^b, heteroaryl substituted with
0-2 R^b, (heteroaryl)C₁₋₄ alkyl substituted with 0-2 R^b,
15 heterocyclyl substituted with 0-2 R^b, and
(heterocyclyl)C₁₋₄ alkyl substituted with 0-2 R^b;

20 R^b is independently selected at each occurrence from C₁₋₄
alkyl, -NO₂, halo, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -
NR⁸C(O)OR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6b}R^{7a}, -NR⁸CONR^{6b}R^{7a}, -
C(O)OR^{10a}, SH, and -S(O)_nR¹²;

25 R^{6b} is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

R⁷ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)methyl;

30 R^{7a} is independently selected at each occurrence from H,
C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl,
(C₃₋₆ cycloalkyl)methyl, and C₁₋₄ haloalkyl;

alternatively, NR^{6a}R^{7a} is selected from piperidine,
pyrrolidine, morpholine, thiomorpholine,
35 thiomorpholine-oxide, and thiomorpholine-dioxide, and
is substituted with 0-1 R^e;

R^e is C₁₋₄ alkyl;

alternatively, NR^{6a}R^{7a} is piperazine or N-methylpiperazine,
and is substituted with 0-1 R^f;

5 R^f is selected from C₁₋₄ alkyl, C(O)C₁₋₄ alkyl, C(O)benzyl,
C(O)OC₁₋₄ alkyl, C(O)O-benzyl, SO₂-C₁₋₄ alkyl, SO₂-
benzyl, and SO₂-phenyl;

10 R⁸ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

R⁹ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

15 R¹⁰ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, and
(C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

20 R^{10a} is independently selected at each occurrence from H,
C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl,
(C₃₋₆ cycloalkyl)C₁₋₄ alkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, and
C₁₋₄ haloalkyl;

25 R^{10b} is independently selected at each occurrence from H,
C₁₋₆ alkyl, C₂₋₄ alkenyl, phenyl, benzyl, C₃₋₆
cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, (C₁₋₄ alkoxy)C₁₋₄
alkyl, and C₁₋₄ haloalkyl;

30 R¹¹ is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, and C₃₋₆ cycloalkyl;

35 R¹² is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl,
C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, aryl, and
(aryl)C₁₋₄ alkyl;

R¹³ is independently at each occurrence selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

5 R^{13a} is independently at each occurrence selected from C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

10 aryl is independently at each occurrence selected from phenyl and naphthyl, and is substituted with 0-3 R⁹;

R⁹ is independently at each occurrence selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halo, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, SH, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -
15 OC(O)R¹⁷, -NR⁸COR^{16a}, -NR⁸CONR^{6a}R^{7a}, -NR⁸C(O)OR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};

20 R¹⁵ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl, benzyl, (phenyl)C₁₋₄ alkyl and (naphthyl)C₁₋₄ alkyl;

25 R¹⁶ is independently at each occurrence selected from H, C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

30 R^{16a} is independently at each occurrence selected from C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₄ alkyl;

35 R¹⁷ is independently at each occurrence selected from C₁₋₄ alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl, C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl, benzyl, (phenyl)C₁₋₄ alkyl, (naphthyl)C₁₋₄ alkyl, heteroaryl and (heteroaryl)C₁₋₄ alkyl;

heteroaryl is independently at each occurrence selected from pyridyl, pyrimidinyl, triazinyl, furanyl, quinolinyl,

isoquinolinyl, thienyl, imidazolyl, thiazolyl, indolyl,
pyrrolyl, oxazolyl, benzofuranyl, benzothienyl,
benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl,
tetrazolyl, and indazolyl, and is substituted with 0-3
5 R^b;

heterocyclyl is saturated or partially saturated heteroaryl,
substituted with 0-3 R^b; and

10 R^b is independently at each occurrence selected from
C₁₋₆ alkyl, C₃₋₆ cycloalkyl, halo, C₁₋₄ haloalkyl, cyano,
nitro, -OR^{10a}, SH, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -
OC(O)R¹⁸, -NR⁸COR^{16a}, -NR⁸CONR^{6a}R^{7a}, -NR⁸CO₂R^{16a}, -
NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};

15 R¹⁸ is independently selected at each occurrence from
C₁₋₆ alkyl, C₃₋₆ cycloalkyl, phenyl and benzyl; and

n is independently at each occurrence selected from 0, 1 and
20 2.

2. A compound according to Claim 1, wherein:

25 R is selected from phenyl and pyridyl, and is substituted
with 0-5 R³ groups;

R¹ is selected from H, CH₃, CH₂CH₃, Cl, F, Br, -CN, CF₃,
NH₂, NH(CH₃), N(CH₃)₂, OH, OCH₃, SH, SCH₃, and
30 S(O)₂CH₃;

R² is NR^{6a}R^{7a}; and,

35 R³ is independently selected at each occurrence from
C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl
substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with
0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a,
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, -

NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹².

5 3. A compound according to Claim 2, wherein:

R is selected from phenyl and pyridyl, and is substituted with 1-3 R³ groups;

10 R¹ is selected from H, CH₃, CH₂CH₃, Cl, and F;

R³ is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, F, Cl, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR¹⁰, and -S(O)_nR¹².

20 R^a is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, F, Cl, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6a}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²;

25 R^{6a} is independently selected at each occurrence from C₁₋₆ alkyl substituted with 0-2 R^b, C₂₋₆ alkenyl substituted with 0-2 R^b, C₂₋₆ alkynyl substituted with 0-2 R^b, C₃₋₆ cycloalkyl substituted with 0-2 R^b, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl substituted with 0-2 R^b, (aryl)C₁₋₄ alkyl substituted with 0-2 R^b, (heteroaryl)C₁₋₄ alkyl substituted with 0-2 R^b, and (heterocyclyl)C₁₋₄ alkyl substituted with 0-2 R^b;

35 R^b is independently selected at each occurrence from C₁₋₄ alkyl, -NO₂, F, Cl, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -COR¹³, -OR^{10a}, -CONR^{6b}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²; and,

R^{7a} is independently selected at each occurrence from H, C₁₋₆ alkyl, benzyl, and (C₃₋₆ cycloalkyl)methyl.

4. A compound according to Claim 3, wherein:

5 R¹ is CH₃;

R³ is independently selected at each occurrence from
C₁₋₆ alkyl substituted with 0-2 R^a, C₂₋₆ alkenyl
substituted with 0-2 R^a, C₂₋₆ alkynyl substituted with
10 0-2 R^a, C₃₋₆ cycloalkyl substituted with 0-2 R^a,
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl substituted with 0-2 R^a, -
NO₂, halo, -CN, C₁₋₄ haloalkyl, -NR⁶R⁷, -NR⁸COR^{13a}, -
COR¹³, -OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹²;

15 R^a is independently selected at each occurrence from C₁₋₄
alkyl, -NO₂, F, Cl, -CN, -NR⁶R⁷, -NR⁸COR^{13a}, -COR¹³, -
OR^{10a}, -CONR⁶R⁷, -C(O)OR^{10a}, and -S(O)_nR¹²;

R^{6a} is independently selected at each occurrence from
20 C₂₋₄ alkyl substituted with 0-2 R^b, (C₃₋₅ cycloalkyl)C₁₋₂
alkyl substituted with 0-2 groups selected from CH₃O
and CH₃CH₂O, (aryl)C₁₋₂ alkyl substituted with 0-2 R^b,
(heteroaryl)C₁₋₂ alkyl substituted with 0-2 R^b, and
(heterocyclyl)C₁₋₂ alkyl substituted with 0-2 R^b;

25 R^b is independently selected at each occurrence from C₁₋₄
alkyl, -NO₂, F, Cl, -CN, -NR^{6b}R^{7a}, -NR⁸COR^{13a}, -COR¹³, -
OR^{10a}, -CONR^{6b}R^{7a}, -C(O)OR^{10a}, and -S(O)_nR¹²;

30 R⁸ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

R⁹ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₃₋₆ cycloalkyl and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

35 R¹⁰ is independently at each occurrence selected from H, C₁₋₄
alkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, and
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

- R^{10a} is independently selected at each occurrence from H,
C₁₋₆ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl,
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl, (C₁₋₄ alkoxy)C₁₋₂ alkyl, and
5 C₁₋₄ haloalkyl;
- R^{10b} is independently selected at each occurrence from H,
C₁₋₆ alkyl, C₂₋₄ alkenyl, phenyl, benzyl, C₃₋₆
cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl, (C₁₋₄ alkoxy)C₁₋₂
10 alkyl, and C₁₋₄ haloalkyl;
- R¹¹ is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, and C₃₋₆ cycloalkyl;
- 15 R¹² is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₂ alkyl,
C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₂ alkyl, aryl, and
(aryl)C₁₋₂ alkyl;
- 20 R^{13a} is independently at each occurrence selected from C₁₋₆
alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and
(C₃₋₆ cycloalkyl)C₁₋₂ alkyl;
- aryl is independently at each occurrence phenyl substituted
25 with 0-3 R⁹;
- R⁹ is independently at each occurrence selected from
C₁₋₄ alkyl, C₃₋₆ cycloalkyl, F, Cl, C₁₋₄ haloalkyl,
cyano, nitro, -OR^{10a}, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -
30 NR⁸COR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a};
- R¹⁵ is independently at each occurrence selected from C₁₋₄
alkyl, C₁₋₄ haloalkyl, (C₁₋₄ alkoxy)C₁₋₄ alkyl,
C₃₋₆ cycloalkyl, (C₃₋₆ cycloalkyl)C₁₋₄ alkyl, phenyl,
35 benzyl, and (phenyl)C₁₋₂ alkyl;

R¹⁶ is independently at each occurrence selected from H, C₁₋₄ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

5 R^{16a} is independently at each occurrence selected from C₁₋₄ alkyl, phenyl, benzyl, C₃₋₆ cycloalkyl, and (C₃₋₆ cycloalkyl)C₁₋₂ alkyl;

10 heteroaryl is independently at each occurrence selected from pyridyl, pyrimidinyl, furanyl, imidazolyl, thiazolyl, pyrrolyl, oxazolyl, isoxazolyl, pyrazolyl, triazolyl, and tetrazolyl, and is substituted with 0-3 R^h;

15 heterocyclyl is saturated or partially saturated heteroaryl, substituted with 0-3 R^h; and,

R^h is independently at each occurrence selected from C₁₋₆ alkyl, C₃₋₆ cycloalkyl, F, Cl, C₁₋₄ haloalkyl, cyano, nitro, -OR^{10a}, -S(O)_nR¹⁵, -COR¹⁶, -C(O)OR¹⁶, -NR⁸COR^{16a}, -NR^{6a}R^{7a}, and -CONR^{6a}R^{7a}.

20

5. A compound according to Claim 1, wherein the compound is selected from the group:

- 25 3-(2,4-dichlorophenyl)-5-methyl-7-(3-pentyl)amino-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dichlorophenyl)-5-methyl-7-(methoxybut-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dichlorophenyl)-5-methyl-7-(1,3-bis(methoxy)prop-2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dichlorophenyl)-5-methyl-7-(di(methoxyethyl)amino)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-butyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-propyl)amino-isoxazolo[4,5-d]pyrimidine;

- 3-(2,4-dichlorophenyl)-5-methyl-7-diethylamino-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dichlorophenyl)-5-methyl-7-dipropylamino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dichlorophenyl)-5-methyl-7-(N-propyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-butyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dichlorophenyl)-5-methyl-
7-di(cyclopropylmethyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dichlorophenyl)-5-methyl-7-(N-ethyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-propyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dichlorophenyl)-5-methyl-7-(N-butyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dichlorophenyl)-5-methyl-7-dicyclopropylamino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(N-cyclopropyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbut-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpent-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dichlorophenyl)-5-methyl-7-((1,3-dicyclopropyl)prop-
2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpropyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbutyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4-dichlorophenyl)-5-methyl-7-
(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-
(1,2-dicyclopropyl)ethyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 55

- 3-(2,4-dichlorophenyl)-5-methyl-7-(1-hydroxy-but-2-yl)amino-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dichlorophenyl)-5-methyl-7-(morpholino)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(3-pentoxy)-
isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dichlorophenyl)-5-methyl-7-(methoxybut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dichlorophenyl)-5-methyl-7-(cyclopropylpropoxy)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dichlorophenyl)-5-methyl-7-(dicyclopropylmethoxy)-
isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dichlorophenyl)-5-methyl-7-(2-methoxy-1-
cyclopropylethoxy)-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4,6-trimethylphenyl)-5-methyl-7-(3-pentyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(methoxybut-2-yl)amino-
isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4,6-trimethylphenyl)-5-methyl-7-(1,3-bis(methoxy)prop-
2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-
35 (di(methoxyethyl)amino)-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-butyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-
propyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-diethylamino-
isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4,6-trimethylphenyl)-5-methyl-7-dipropylamino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-
50 cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-propyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 55 3-(2,4,6-trimethylphenyl)-5-methyl-7(N-butyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;

- 3-(2,4,6-trimethylphenyl)-5-methyl-
7-di(cyclopropylmethyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-ethyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-propyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-butyl-N-
cyclopropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4,6-trimethylphenyl)-5-methyl-7-dicyclopropylamino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(N-cyclopropyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylbut-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylpent-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4,6-trimethylphenyl)-5-methyl-7-((1,3-
dicyclopropyl)prop-2-yl)amino-
isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4,6-trimethylphenyl)-5-methyl-7-
(cyclopropylpropyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-
(cyclopropylbutyl)amino-isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4,6-trimethylphenyl)-5-methyl-7-
(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-((1,2-
dicyclopropyl)ethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4,6-trimethylphenyl)-5-methyl-7-(1-hydroxy-but-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4,6-trimethylphenyl)-5-methyl-7-(morpholino)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(3-pentoxy)-
isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4,6-trimethylphenyl)-5-methyl-7-(methoxybut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 55 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;

- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(cyclopropylpropoxy)-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4,6-trimethylphenyl)-5-methyl-7-(dicyclopropylmethoxy)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4,6-trimethylphenyl)-5-methyl-7-(2-methoxy-1-
cyclopropylethoxy)-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dimethylphenyl)-5-methyl-7-(3-pentyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(methoxybut-2-yl) amino-
isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dimethylphenyl)-5-methyl-7-(1,3-bis(methoxy) prop-2-
yl) amino-isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dimethylphenyl)-5-methyl-7-(di(methoxyethyl) amino)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-butyl) amino-
isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-propyl) amino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-diethylamino-
isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dimethylphenyl)-5-methyl-7-dipropylamino-
isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-
cyclopropylmethyl) amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-propyl-N-
cyclopropylmethyl) amino-isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dimethylphenyl)-5-methyl-7(N-butyl-N-
cyclopropylmethyl) amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-
7-di(cyclopropylmethyl) amino-
isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dimethylphenyl)-5-methyl-7-(N-ethyl-N-
cyclopropyl) amino-isoxazolo[4,5-d]pyrimidine;
- 50 3-(2,4-dimethylphenyl)-5-methyl-7-(N-propyl-N-
cyclopropyl) amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(N-butyl-N-
cyclopropyl) amino-isoxazolo[4,5-d]pyrimidine;
- 55

- 3-(2,4-dimethylphenyl)-5-methyl-7-dicyclopropylamino-
isoxazolo[4,5-d]pyrimidine;
- 5 3-(2,4-dimethylphenyl)-5-methyl-7-(N-cyclopropyl-N-
cyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbut-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 10 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpent-2-
yl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-((1,3-dicyclopropyl)prop-
2-yl)amino-isoxazolo[4,5-d]pyrimidine;
- 15 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpropyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbutyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 20 3-(2,4-dimethylphenyl)-5-methyl-7-
(dicyclopropylmethyl)amino-isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-
((1,2-dicyclopropyl)ethyl)amino-
isoxazolo[4,5-d]pyrimidine;
- 25 3-(2,4-dimethylphenyl)-5-methyl-7-
(1-hydroxy-but-2-yl)amino-
isoxazolo[4,5-d]pyrimidine;
- 30 3-(2,4-dimethylphenyl)-5-methyl-7-(morpholino)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(3-pentoxo)-
isoxazolo[4,5-d]pyrimidine;
- 35 3-(2,4-dimethylphenyl)-5-methyl-7-(methoxybut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 40 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylbut-2-oxy)-
isoxazolo[4,5-d]pyrimidine;
- 3-(2,4-dimethylphenyl)-5-methyl-7-(cyclopropylpropoxy)-
isoxazolo[4,5-d]pyrimidine;
- 45 3-(2,4-dimethylphenyl)-5-methyl-7-(dicyclopropylmethoxy)-
isoxazolo[4,5-d]pyrimidine; and,
- 50 3-(2,4-dimethylphenyl)-5-methyl-7-(2-methoxy-1-
cyclopropylethoxy)isoxazolo[4,5-d]pyrimidine;

or a pharmaceutically acceptable salt form thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/18707

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D498/04 A61K31/505 //(C07D498/04,261:00,239:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 08847 A (PFIZER ;CHEN YUHPYNG LIANG (US)) 5 March 1998 (1998-03-05) cited in the application see claim 1 and page 8, Formula I-S	1-5
X	CHEMICAL ABSTRACTS, vol. 66, no. 13, 27 March 1967 (1967-03-27) Columbus, Ohio, US; abstract no. 55453, DESIMONI, GIOVANNI ET AL: "Polynuclear isoxazole types. I. Isoxazolo[4,5-d]pyrimidines" XP002124218 RN: 14258-83-4, 14258-84-5, 14258-85-6, 14258- 86-7, 14258-87-8, 14258-88-9, 14258-89-0, 5993 -86-2 abstract & TETRAHEDRON (1967), 23(2), 675-80 , -/-	1,2
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
Date of the actual completion of the international search 29 November 1999		Date of mailing of the international search report 14/12/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Steendijk, M

INTERNATIONAL SEARCH REPORT

Inter: nat Application No
PCT/US 99/18707

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 122, no. 13, 27 March 1995 (1995-03-27) Columbus, Ohio, US; abstract no. 160584, LAURENT, SOPHIE ET AL: "Synthesis of isoxazolo'4,5-d!pyrimidinones, and their isomerization into oxazolo'4,5-d!pyrimidinones on flash vacuum pyrolysis" XP002124219 RN:160687-68-3 abstract & BULL. SOC. CHIM. BELG. (1994), 103(5-6), 181-4 ,	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. nat. Application No

PCT/US 99/18707

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9808847 A	05-03-1998	AU 3456197 A	19-03-1998
		CN 1227554 A	01-09-1999
		EP 0923582 A	23-06-1999
		HR 970454 A	31-08-1998
		NO 990927 A	26-02-1999
		PL 332040 A	16-08-1999